```
2000:36379
                CAPLUS
\Delta N
DN
   133:69901
    Entered STM: 17 Jan 2000
MD.
    Nodel dendrons and dendrimers incorporating diphenylamino-substituted .
TI
     diphenylpolyene and PPV-cligomer moieties for NLO applications
    Astanorin, Kimba: Spangler, Charles W.; Reeves, Benjamin
AU
     Dep. Chem. Biochem., Montana State Univ., Bozeman, MT, USA
CS
     Proceedings of SPIE-The International Society for Optical Engineering
SO
     (1999), 3796 (Organic Wonlinear Optical Materials), 170-177
     CODEN: PSISDG; ISSN: 0277-786X
     SPIN-The International Society for Optical Engineering
233
DT
    Journal.
    Boglish
1.2
೦೦
     35-8 (Chemistry of Synchetic Righ Polymers)
     Section cross-reference(s): 36, 73
    The synthesis and characterization are described of diphenylamino-
XX.
     substituted diphenylpolyene and poly(p-phenylenevinylene)s as two-photon
     absorbers, photoluminescent materials suitable for org. light-emitting
     diodes, and as dendrimers capable of 3D charge delocalization and
     exceptionally large third order hyperpolarizability. Bis-
     (diphenylamino) diphenylpolyenes form exceptionally stable, highly
     absorbing bipolaronic dications in soln, and thin film. Replacement of
     one diphonylamino substituent with a N-(hydroxyethyl), N-sthylaminophenyl
     sciety yields a polyene that also forms stable bipolarons which are
     intensely fluorescent. These chromophores are easily attached to either a
     PMMA backbone or to 3,5-dihydroxybenzyl alc. to yield functionalized
     dendrons capable of attachment to various core mols. Diphenylamino-
     substituted FFV oligomers can also be obtained with similar functionality.
     These materials possess large two-photom cross-sections and display
     optical limiting for manosecond pulses with peak activity extending into
     the visible portion of the spectrum.
ST
     dendron diphenylamino diphenylpolyene prepn monlinear optical property;
     polyphenylenevinylene diphenylasino substituent photon absorber: optical
     limiting dendrimer diphenylamine diphenylpolyene
\mathfrak{T}^{m}
     Poly(arylenes)kenylenes)
     RL: FNG (Preparation, unclassified); PRP (Properties); PRSP (Preparation)
        (dendritie; prepn. of model dendrons and dendrimers incorporating
        diphenylamino-diphenylpolyene and FPV-oligomer and luminescence and
        hyperpolarizability of compds. for WLO applications)
IT
     Chemical chains
        (hyperbranched; preph. of model dendrons and dendrimers incorporating
        diphenylamino-diphenylpolyene and PFV-oligomer and luminescence and
        hyperpolarizability of compds. for NLO applications)
TT
    Sipolaron
    Fluorescence
    Luminescence, electroluminescence
    Nonlinear optical materials
    Optical hyperpolarizability
     Optical limiting
        (prepa. of model dendross and dendrimers incorporating
        diphonylamino-diphenylpolyene and PPV-oligomer and luminescence and
        hyperpolarizability of compds. for NLO applications)
TT
    Dendritic polymers
```

RL: PNU (Preparation, unclassified); RCT (Reactant); PREP (Preparation); FACT (Reactant or reagent)

RL: PNU (Preparation, unclassified); FRP (Properties); PREP (Preparation)

diphenylamino-diphenylpolyene and PFV-oligomer and luminescence and

(prepn. of model dendrons and dendrimers incorporating

hyperpolarizability of compds. for NLO applications)

TT

281655-29-6P

(convergent dendron; prepn. of model dendrons and dendrimers incorporating diphenylamino-diphenylpolyene and PPV-oligomer and luminescence and hyperpolarizability of compds. for NLO applications)

IT 281655-31-0P

RL: PNU (Preparation, unclassified); PRP (Properties); PREP (Preparation) (dendrimer; preparation) diphenylamino-diphenylpolyene and PPV-oligomor and luminescence and hyperpolarizability of compds. for NLO applications)

IT 134061-63-5P

RL: PNU (Preparation, unclassified); PRP (Properties); PRBP (Preparation) (prepn. of model dendrons and dendrimers incorporating diphenylamino-diphenylpolyens and PPV-oligomer and luminescence and hyperpolarizability of compde. for NLO applications):

TT 281655-28-5P, 4-Diphenylamino-4'-(N-ethyl-N-(2-hydroxyethyl))stilbene RL: PNU (Preparation, unclassified); RCT (Reactant); PREP (Preparation); RACT (Reactant or respent)

(prepn. of model dendrons and dendrimers incorporating diphenylamino-diphenylpolyene and PPV-oligomer and luminescence and hyperpolarizability of compds. for NLO applications)

IT 55055-42-2, 4-(Diphenylamine)-4'-[4-(diphenylamine)styryl]stilbene RL: PRF (Properties)

(prepn. of model dendrons and dendrimers incorporating diphenylamino-diphenylpolyene and PPV-oligomer and luminescence and hyperpolarizability of compds. for NLO applications)

'T 80-05-7, reactions 29654-55-5, 3,5-Dihydroxybenzyl alcohol
RL: RCT (Reactant); RACT (Reactant or reagent)
(prepn. of model dendrons and dendrimers incorporating
diphenylamino-diphenylpolyene and PPV-oligomer and luminescence and
hyperpolarizability of compds. for NLO applications)

RE.(NT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD RE

- (1) Anom; private communication from L S Sapochak
- (2) Anon; private communication from L V Natarajan
- (3) Ashworth, K; Mat Res Soc Symp Proc, to be published 1999
- (4) Ehrlich, J; Mat Res Soc Symp Proc 1997, V479, P9 <u>CAPLUS</u>
- (5) Hawker, C; Macromol 1990, V23, P4726 CAPLUS
- (6) Reeves, B; Honore Thesis, Montana State University 1999
- (7) Spangler, C: Handbook of Conductive Molecules and Polymers: Vol 2. Conductive Polymers: Synthesis and Electrical Properties 1997, P389
- (8) Spangler, C: J Chem Soc Perkin Trans 1 1989, P151 CAPLUS
- (8) Spangler, C; J Chem Soc Perkin Trans 1 1991, P799 CAPLUS
- (10) Spangler, C; J Chem Soc Perkin Trans 1 1995, P715 CAPLOS
- (11) Spangler, C; Mat Res Soc Symp Proc 1998, V488, F283 CAPLUS
- (12) Wooley, K; J Chem Soc Perkin Trans 1 1991, F1059 CAPLUS